



## 10 supozitorija sa efervescentnim laksativnim dejstvom sa Kamilicom

\* Supozitorije za ne posredan i efikasan, prirodan tretman protiv konstipacije.

\* Sadrži kamilicu da pomogne u olakšanju osećaja nepotpune defekacije

\* Odlikuje se restauracijom mehanizma za defekaciju



### Konstipacija

Konstipacija se uglavnom karakteriše smanjenjem motilitetom creva i karakteriše se tvrdom i suvom stolicom.

Najčešće se javlja zbog neuobičajeno male pokretljivosti creva, što dovodi do akumulacije stolice u donjem delu. Frekvenca pokretljivosti ljudskog creva varira od 3 puta dnevno do 3 puta nedeljno. Uobičajeno je da je konstipacija praćena simptomima različitih stepena težine, od kojih su neki anoreksija, tupa glavobolja, iscrpljenost, bol u ledima, abdominalna nadutost, bol u stomaku.

### EVA LAX®

**EVA LAX®** je proizvod u obliku supozitorija za lečenje konstipacije izazvane delimičnom blokadom defekacije na prirodan način. Specijalno je formulisani da olakša kretanje creva u slučaju konstipacije, dok istovremeno smanjuje nadraženost rektalnog sfinktera usled osećaja nepotpune defekacije i umiruje iritaciju rektuma i osećaj pečenja.

**EVA LAX®** supozitorije nemaju farmakološki efekat i ne izazivaju toleranciju.

### Mehanizam dejstva

Nakon aplikacije u rektum, **EVA LAX®** supozitorija se rastvara u roku od nekoliko minuta, stvarajući mehuriće (šumeće dejstvo), koji pritisaku zidove creva, potiskujući stolicu van debelog creva. Istovremeno, polietilenglikol iz rastvorene supozitorije omekšava stolicu, dok kamilica pomaže u smanjenju mogućeg osećaja neprijatnosti usled iritacije creva, zbog njenog umirujućeg dejstva. Efekat supozitorija je momentalan i izaziva pokretanje creva u roku od 5-30 minuta. Mechanizam delovanja proizvoda je fizičko-mehanički bez ikakvog farmakološkog efekta. Preparat se izbacuje nepromjenjen, bez absorpcije u crevima.

### Indikacije

**EVA LAX®** supozitorije su indikovane za lečenje povremene i recidivirane konstipacije na prirodan način. **EVA LAX®** se takođe može koristiti za pripremu pre rektalnog pregleda i

proktoskopije. Zbog svog prirodnog efekta mogu ga koristiti starije osobe, osobe sa problemima pri kretanju, a takođe se može primeniti i kod konstipacije kod putika.

### Neželjeni efekti

**EVA LAX®** supozitorije ne izazivaju neželjene efekte, čak i u slučajevima česte i dugotrajne upotrebe, jer se sastoje od netoksičnih supstanci koje su zastupljene u proizvodnji hrane i lekova opisanih u Evropskoj farmakopeji, kao i u drugim farmakopejama.

### Upotreba

Skinite plastični poklopac, pokvasite supozitoriju vodom (za najbolje rezultate stavite supozitoriju pod tekuću vodu 5-10 sekundi) pre upotrebe i ubacite u rektum. Nakon nanošenja, supozitorija će početi da se rastvara i delovaće odmah. Zadržite supozitoriju u rektumu što je duže moguće (10-30 minuta) dok se krećete, pre defekacije.

Ukoliko patite od hemoroida ili drugih problema u analnoj regiji, preporučuje se podmazivanje supozitorija maslinovim uljem, suncokretnim uljem ili vazelinom pre upotrebe.

### Mere predostrožnosti

- Pogodno za odrasle i decu stariju od 12 godina.
- Za decu do 12 godina konsultujte svog lekara.
- Za upotrebu tokom trudnoće ili dojenja, konsultujte svog lekara.
- Koristite jednu ili više supozitorija za svaki tretman.
- Neželjeni efekati nisu zabeleženi nakon upotrebe **EVA LAX®** supozitorija. Osećaj nepotpune defekacije ili umerenog prelaznog bola ne bi trebao da vas brinu zbog toga što su to simptomi koji se mogu tretirati. Nema poznatih kontraindikacija, čak i u slučaju česte i dugotrajne upotrebe, kao i u prisustvu hemoroida i lokalnih oštećenja.
- Krvarenje ili nemogućnost defekacije nakon upotrebe **EVA LAX®** može značiti ozbiljno stanje. Prekinite upotrebu i konsultujte svog lekara.
- Čuvati na temperaturi od 20-25°C na tamnom mestu, van domaća dece.

#### Proizvođač:

IOULIA AND IRENE TSETI Pharmaceutical Laboratories S.A., "INTERMED S.A."

Ulica: 27 Kalyftaki str, 145 64 Kifissia, GRČKA

Nosilac upisa, uvoznik i distributer za Srbiju: Braf d.o.o., 11080 Beograd - Zemun, Lazara Mamuzića 28

Rešenje broj: 515-02-00851-18-001 od 03.05.2019.

ZA VIŠE INFORMACIJA POSETITE: [www.mojaEVA.rs](http://www.mojaEVA.rs)

Nosilac upisa, uvoznik i distributer za Crnu Goru: Poen d.o.o., 85320 Tivat, Donja Lastva bb

Rješenje broj: 2070/18/876-6115 od 27.11.2018.

ZA VIŠE INFORMACIJA POSETITE: [www.mojaEVA.me](http://www.mojaEVA.me)





## 10 Suppositories with effervescent laxative action with Chamomile

\*Suppositories for the immediate and effective natural confrontation of constipation.

\*Contains chamomile to help relieve tenesmus.

\*Favors the restoration of the defecation mechanism.



### Constipation

Constipation is generally determined by the decrease in frequency of bowel movement and is characterized by the difficult passing of hard and dry stool.

This usually occurs because of an abnormally slow motility of stool passing the colon, causing accumulation of stool at the bottom part. The mobility frequency of human bowel varies from 3 times per day to 3 times per week. Usually, constipation is accompanied by symptoms of various severity rates some of which are anorexia, dull headache, exhaustion, lower back pain, abdominal distention, lower abdominal pain.

### EVA LAX®

**EVA LAX®** is a product in the form of suppositories for the treatment of constipation caused by partial stool blockage, in a natural way.

It has been specially designed to facilitate bowel movement in cases of constipation, while reducing rectal sphincter tenesmus and soothing rectal irritation and burning.

**EVA LAX®** suppositories have no pharmacological effect and do not cause tolerance.

### Action Mechanism

After being inserted in the rectum, the **EVA LAX®** suppository is dissolved within a few minutes, creating microbubbles (effervescent action), which apply pressure to the walls of the intestine, pushing stool out of the large intestine. At the same time, polyethylene glycol from the dissolved suppository softens the stool while chamomile helps reduce possible burning sensation from intestinal irritation, because of its soothing action. The suppository's effect is immediate and provokes bowel movement in 5-30 minutes. The product's action mechanism is purely physico-mechanical without any pharmacological effect. The material is excreted unchanged without being absorbed by the large intestine.

### Indications

**EVA LAX®** suppositories are indicated for the treatment of occasional and relapsing constipation, in a natural way. **EVA LAX®** can also be used for preparation prior to rectal

examination and proctoscopy. Because of its natural effect it can be used by the elderly, by people with mobility problems and also to confront traveler's constipation.

### Adverse effects

**EVA LAX®** suppositories do not cause adverse effects to the body, even in cases of frequent and long-term use, because it is composed of non-toxic substances which are widely used in the manufacture of food and medicines described in the European Pharmacopeia as well as other pharmacopeias.

### Use

Remove the plastic cover, wet the suppository in some water (for best results place the suppository under running water for 5-10 seconds) before its use and insert in the rectum. After its application, the suppository will begin to dissolve and act immediately. Keep the suppository in the rectum for as long as possible (10-30 minutes) while moving, before defecation. If you suffer from hemorrhoids or other anal discomfort it is recommended to lubricate the suppository with olive oil, sunflower oil or Vaseline before placement.

### Precautions

- Suitable for adults and children over the age of 12.
- For children under 12 consult your physician.
- For use during pregnancy or lactation, consult your physician prior to use.
- Use one or more suppositories for each treatment.
- No side effects have been reported from the use of **EVA LAX®** suppositories. A feeling of insufficient defecation or moderate transient pain should not worry you because these are symptoms which can be treated. There are no known contraindications even during frequent and prolonged use and in the presence of hemorrhoids and local cracks.
- Bleeding or failure to defecate after the use of **EVA LAX®** can imply a serious condition. Discontinue use and consult your physician.
- Store at a temperature of 20-25°C in a dark place, out of children's reach.



#### Manufacturer:

IOULIA AND IRENE TSETI Pharmaceutical Laboratories S.A., "INTERMED S.A."

Address: 27 Kalyftaki str, 145 64 Kifissia, GREECE

Registration holder, importer and distributor for Serbia: Braf d.o.o.,

11080 Beograd - Zemun, Lazara Mamuzića 28

Registration number: 515-02-00851-18-001 od 03.05.2019.

FOR MORE INFORMATION PLEASE CONSULT: [www.mojaEVA.rs](http://www.mojaEVA.rs)

Registration holder, importer and distributor for Montenegro:/

Poen d.o.o., 85320 Tivat, Donja Lastva bb

Registration number: 2070/18/876-6115 od 27.11.2018.

FOR MORE INFORMATION PLEASE CONSULT: [www.mojaEVA.me](http://www.mojaEVA.me)

